



Complete mitochondrial genome of the freshwater monogonont rotifer *Brachionus rubens* (Rotifera, Brachionidae)

Beom-Soon Choi, Young Hwan Lee, Jin-Sol Lee, Erick O. Ogello, Hee-Jin Kim, Atsushi Hagiwara & Jae-Seong Lee

To cite this article: Beom-Soon Choi, Young Hwan Lee, Jin-Sol Lee, Erick O. Ogello, Hee-Jin Kim, Atsushi Hagiwara & Jae-Seong Lee (2020) Complete mitochondrial genome of the freshwater monogonont rotifer *Brachionus rubens* (Rotifera, Brachionidae), *Mitochondrial DNA Part B*, 5:1, 5-6, DOI: [10.1080/23802359.2019.1694853](https://doi.org/10.1080/23802359.2019.1694853)

To link to this article: <https://doi.org/10.1080/23802359.2019.1694853>



© 2019 The Author(s). Published by Informa UK Limited, trading as Taylor & Francis Group.



Published online: 10 Dec 2019.



Submit your article to this journal [↗](#)



View related articles [↗](#)



View Crossmark data [↗](#)

Complete mitochondrial genome of the freshwater monogonont rotifer *Brachionus rubens* (Rotifera, Brachionidae)

Beom-Soon Choi^{a*}, Young Hwan Lee^{b*}, Jin-Sol Lee^b, Erick O. Ogello^c, Hee-Jin Kim^d, Atsushi Hagiwara^{d,e} and Jae-Seong Lee^b

^aPhyzen Genomics Institute, Seongnam, South Korea; ^bDepartment of Biological Science, College of Science, Sungkyunkwan University, Suwon, South Korea; ^cDepartment of Fisheries & Natural Resources, Maseno University, Kisumu, Kenya; ^dInstitute of Integrated Science and Technology, Nagasaki University, Nagasaki, Japan; ^eOrganization for Marine Science and Technology, Nagasaki University, Nagasaki, Japan

ABSTRACT

The two complete mitochondrial genomes were sequenced from the freshwater monogonont rotifer *Brachionus rubens*. The genome sequences were 12,041 bp and 13,793 bp in size, and the gene order and contents were identical to those of the freshwater rotifer *B. rubens* China, but were different in three tRNA-Arg, tRNA-Ile, and tRNA-Leu between both *B. rubens* mitochondrial genomes, while *B. calyciflorus* had peculiar gene order in mitochondrial DNA I. Of 12 protein-coding genes (PCGs), one gene (*ND5*) had incomplete stop codons. Furthermore, the start codon of *ND4* and *CO2* gene was ATT, while the start codon of other PCGs was ATG. The base composition of 12 PCGs in *B. rubens* mitogenome showed 22.5% for A, 46.5% for T, 16.3% for C, and 14.7% for G, respectively.

ARTICLE HISTORY

Received 16 October 2019
Accepted 27 October 2019

KEYWORDS



Monogonont rotifer;
complete mitochondrial
genome; *Brachionus rubens*

To date, only a few complete mitochondrial genomes have been published in the freshwater rotifer *Brachionus* sp.; *Brachionus calyciflorus* (Nie et al. 2016) and *Brachionus rubens* China (GenBank KJ489417 and KJ489418), while several marine rotifer *Brachionus* sp. mitogenomes were reported (Suga et al. 2008 for *B. plicatilis*; Hwang et al. 2014 for *Brachionus koreanus*, and Kim et al. 2017 for *Brachionus rotundiformis*). *Brachionus rubens* is one of the major cosmopolitan freshwater rotifers and was considered as a model for environmental toxicology in response to toxic cyanobacterium (Geng et al. 2006; Geng and Xie 2008; Pérez-Morales et al. 2014), temperature and metal (Azua-García et al. 2006; Montúfar-Meléndez et al. 2007), suggesting that this species can be an important member of freshwater zooplankton community as a sentinel species for ecotoxicology. The analysis of *B. rubens* mitochondrial genome is important to identify field-sampled and laboratory stocks. In this study, we identified two complete mitochondrial genomes of the monogonont rotifer *B. rubens* Japan to better understand the phylogenetic placement of the freshwater rotifer *Brachionus* clade.

The adult *B. rubens* were collected from the freshwater catchment area at Urakami, Nagasaki in Japan (32°79'90.72"N, 129°86'55.32"E) in August 2004 and maintained at the Laboratory of Professor Atsushi Hagiwara, Nagasaki University in Japan. The type was deposited in the ichthyological collection of the Faculty of Fisheries, Nagasaki University (FFNU) under the accession no. FFNU-Rot-0003. We sequenced 300 bp, 500 bp and 800 bp paired end library

of *B. rubens* from whole body genomic DNA using the Illumina HiSeq 2500 platform (GenomeAnalyzer, Illumina, San Diego, CA). *De novo* assembly was conducted by SPAdes (version 3.13.0) (<http://cab.spbu.ru/software/spades/>) with K-mer auto. Of the assembled *B. rubens* 270,986 contigs with Newbler (version 2.9; identity 97) (<http://www.454.com>), seven mitochondrial contigs were obtained. After a manual curation of seven contigs with Consed (version 19.0) (<http://www.phrap.org/consed/consed.html>), two contigs were obtained to the mitochondrial DNA of *B. rubens*.

The complete mitochondrial genomes of *B. rubens* were 12,041 bp (mitochondrial DNA I; GenBank no. MN256531) and 13,793 bp (mitochondrial DNA II; GenBank no. MN256532) in size. The direction of 12 protein-coding genes (PCGs) of *B. rubens* Japan was identical to those of *B. rubens* China of the genus *Brachionus*, including the presence of nearly identical non-coding region (Identities: 5651/5731 = 98.25%) (Suga et al. 2008; Hwang et al. 2013). Of 12 PCGs, one gene (*ND5*) had incomplete stop codons. Furthermore, the start codon of *ND4* and *CO2* gene was ATT, while the start codon of other PCGs was ATG. The base composition of 12 PCGs in *B. rubens* Japan mitogenome showed 22.5% for A, 46.5% for T, 16.3% for C, and 14.7% for G, respectively. The mitochondrial genome A + T base composition (69.0%) of 12 PCGs was higher than G + C (31.0%), while the complete mitochondrial genome A + T base composition (67.5%) was higher than G + C (32.5%).

CONTACT Jae-Seong Lee  jslee2@skku.edu  Department of Biological Science, College of Science, Sungkyunkwan University, Suwon 16419, South Korea
*These authors equally contributed to this work.

© 2019 The Author(s). Published by Informa UK Limited, trading as Taylor & Francis Group.
This is an Open Access article distributed under the terms of the Creative Commons Attribution License (<http://creativecommons.org/licenses/by/4.0/>), which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.




Figure 1. Phylogenetic analysis of the rotifer *Brachionus rubens* Japan mitochondrial DNA. We conducted a comparison of two mitochondrial DNA genes (CO1 and Cytb) of Acoela and Rotifera. Two mitochondrial DNA genes CO1 and Cytb were aligned by ClustalW. Maximum likelihood (ML) analysis was performed by Raxml 8.2.8 (<http://sco.h-its.org/exelixis/software.html>) with GTR + γ + I nucleotide substitution model. The rapid bootstrap analysis was conducted with 1,000 replications with 48 threads running in parallel. The Acoela served as an outgroup. Ln=-5316.105. Modified from Choi et al. (2019).

The placement of *B. rubens* Japan in the genus *Brachionus* with CO1 and Cytb was shown in Figure 1. *B. angularis* was clustered closely to *B. rubens* China. The gene order and contents of 12 PGCs were identical to those of the freshwater rotifer *B. rubens* China. However, interestingly, the order of tRNAs of three tRNA-Arg, tRNA-Ile, and tRNA-Leu was different between both *B. rubens* mitochondrial genomes, while the freshwater rotifer *B. calyciflorus* had peculiar gene order in mitochondrial DNA I. This indicates that the rearrangement of tRNAs is likely occurring in sporadic manner in the genus *Brachionus* (Hwang et al. 2014).

Disclosure statement

No potential conflict of interest was reported by the authors.

Funding

This work was supported by the Collaborative Genome Program of the Korea Institute of Marine Science and Technology Promotion (KIMST) funded by the Ministry of Oceans and Fisheries (MOF) [No. 20180430]. This work was also supported by a short-term invitation fellowship of the Japan Society of the Promotion of Science [JSPS ID: S19056] to Jae-Seong Lee.

References

Azuara-García R, Sarma SS, Nandini S. 2006. The combined effects of zinc and alga on the life table demography of *Anuraeopsis fissa* and *Brachionus rubens* (Rotifera). *J Environ Sci Health A*. 41(4):559–572.

- Choi B-S, Lee YH, Hagiwara A, Lee J-S. 2019. Complete mitochondrial genome of the freshwater monogonont rotifer *Brachionus calyciflorus* (Rotifera, Brachionidae). *Mitochondrial DNA B*. 4(2):3593–3595.
- Geng H, Xie P. 2008. Experimental studies on the effects of toxic *Microcystis aeruginosa* PCC7820 on the survival and reproduction of two freshwater rotifers *Brachionus calyciflorus* and *Brachionus rubens*. *Ecotoxicology*. 17(8):709–715.
- Geng H, Xie P, Xu J. 2006. Effect of toxic *Microcystis aeruginosa* PCC7820 in combination with a green alga on the experimental population of *Brachionus calyciflorus* and *B. rubens*. *Bull Environ Contam Toxicol*. 76(6):963–969.
- Hwang D-S, Dahms H-U, Park HG, Lee J-S. 2013. A new intertidal *Brachionus* and intrageneric phylogenetic relationship among *Brachionus* as revealed by allometry and CO1-ITS1 gene analysis. *Zool Stud*. 52(1):13.
- Hwang D-S, Suga K, Sakakura Y, Hagiwara A, Park HG, Rhee J-S, Lee J-S. 2014. Complete mitochondrial genome of the monogonont rotifer, *Brachionus koreanus* (Rotifera: Brachionidae). *Mitochondrial DNA B*. 29–30.
- Kim H-S, Hwang D-S, Kim H-J, Sakakura Y, Hagiwara A, Lee J-S. 2017. Complete mitochondrial genome of the monogonont rotifer *Brachionus rotundiformis* (Rotifera, Brachionidae). *Mitochondrial DNA B*. 2(1):39–40.
- Montúfar-Meléndez AI, Sánchez-Ortíz JR, Sarma SS, Nandini S. 2007. Combined effects of temperature and lead concentration on the competition between the rotifers *Brachionus havanaensis* and *Brachionus rubens* (Rotifera: Brachionidae). *J Environ Sci Health A*. 42(3):393–398.
- Nie, Z-J, Gu, R-B, Du, F-K, Shao, N-L, Xu, P, Xu, G-C. 2016. Monogonont rotifer, *Brachionus calyciflorus*, possesses exceptionally large, fragmented mitogenome. *PLoS One* 11(12):e0168263.
- Pérez-Morales A, Sarma SS, Nandini S. 2014. Feeding and filtration rates of zooplankton (rotifers and cladocerans) fed toxic cyanobacterium (*Microcystis aeruginosa*). *J Environ Biol*. 35(6):1013–1020.
- Suga K, Mark Welch DB, Tanaka Y, Sakakura Y, Hagiwara A. 2008. Two circular chromosomes of unequal copy number make up the mitochondrial genome of the rotifer *Brachionus plicatilis*. *Mol Biol Evol*. 25(6): 1129–1137.